

Video Education Increases Patient Attainment of Target Phosphorus Levels

S Boyd, MSW, LCSW;¹ T Christopher Bond, PhD;¹ T Zimmerman, RD;¹ K Parker;¹ K Spach, PhD;¹ D Dunn, MSW¹

¹DaVita Inc., Denver, CO, USA

Introduction

The value of video education in the dialysis setting is unknown. We assessed the acceptance of a video education project and its effectiveness in improving the percent of patients with phosphorus levels at or below the recommended target (≤ 5.5 mg/dL).

Methodology

- Eleven of 13 dialysis centers (n = 13 to 141) in one region of a large dialysis organization (LDO) participated in the video education project.
- Lab results for patients were compared for the 6 months before the intervention (8/09-1/10) to the 3 months after the intervention (4/10-6/10).
- A mixed linear model (PROC MIXED, SAS 9.2) was employed with a repeated measure by center over the 3 time periods (before, during, and after intervention). The first and third periods were contrasted.
- The analysis was weighted by the number of patients at each center due to the wide range of center sizes (13 to 141 patients per center).
- The 2 non-participating centers are provided for reference only and cannot be used for direct comparison.

Results

Video Education Project

- One video was shown for eight consecutive weeks beginning February 1, 2010 to in-center hemodialysis patients while the patients were receiving their treatments.
- Patients were given a hand-out summarizing the main topic of each video.
- Watching the videos was encouraged but voluntary.
- Videos offered (by week):
- The Dialysis Experience: Parts 1 & 2
- Sharon's Place
- Adequate Dialysis
- Living Well on Dialysis
- Diet and Dialysis
- Delicate Balance
- DaVita Kitchen
- Go Fistula: A Patient's CathAway

Figure 1. Percentage of Patients Achieving ≤ 5.5 mg/dL Phosphorus at Participating Centers




Table 1. Percent of Patients Achieving ≤ 5.5 mg/dL Phosphorus

Percent of Patients Achieving ≤ 5.5 mg/dL Phosphorus							
	Mean # of pts with monthly P result	Before program (8/09–1/10)	During program (2/10–3/10)	After program (4/10–6/10)	Change (After–Before)		
Participatin	g Center						
A	141	65.7	68.4	72.5	6.8		
В	57	74.4	76.1	72.2	-2.2		
C	79	75.9	77.7	77.6	1.7		
D	13	55.8	62.5	82.1	26.2		
Ε	33	68.5	58.0	61.4	-7.2		
F	24	67.5	64.0	76.4	9.0		
G	21	75.7	71.9	66.7	-9.1		
Н	38	75.0	73.0	72.3	-2.8		
	40	65.9	72.8	73.5	7.6		
J	53	71.8	69.3	76.0	4.2		
K	36	72.9	67.1	70.3	-2.7		
Center-level Mean 49		69.9	69.2	72.8	2.9		
Patient-level Mean		70.3	70.4	73.0	2.7		

Non-participati Center					
Non-A	13	79.6	88.7	70.3	-9.3
Non-B	72	82.7	77.9	81.7	-1.0
Center-level Mean 42		81.2	83.3	76.0	-5.2
Patient-level Me	an	82.4	79.6	80.1	-2.3

- The percentage of patients achieving target levels for phosphorous (≤ 5.5 mg/dL) was higher after participation in a video education program intended to improve patient phosphorus levels (Table 1 and Figure 1).
- Of the over 300 patients who completed a post-video questionnaire:
- 79% indicated that videos increased their overall knowledge of dialysis
- 80% wanted more video education in the future
- 87% reported the video handouts reinforced what they had learned

Conclusions

- The percentage of patients achieving ≤ 5.5 mg/dL phosphorus after a video education program increased 2.9%, which was marginally significant (p = 0.059).
- The use of video education appears to be an effective way to improve phosphorus levels in dialysis patients.
- Due to resources requirements for such programs, corroboration of effectiveness with more rigorous study designs would be desirable.

Acknowledgements

Our sincere appreciation to the teammates in over 1600 DaVita clinics who work every day not only to take care of patients but also to ensure the extensive data collection on which our work is based. We thank DaVita Clinical Research® (DCR®) for the support in preparing this poster. DCR is committed to advancing the knowledge and practice of kidney care.

Correspondence: shaun.boyd@davita.com

American Nephrology Nurses' Association 43rd National Symposium, April 29–May 2, 2012; Orlando, FL